

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Fig. 1. Testicular ACE [2]. Ribbon model of tACE with an inhibitor, lisinopril, bound to the active site (space-filling red, blue and green atoms). The lid formed by helices $\alpha 1$ and $\alpha 2$ is indicated as a thin blue wire above the active site. The 7 evolutionarily conserved cysteines are indicated in yellow. The 2 cystines with their associated short β -sheets are visible. The cystine on the right of the figure links helix $\alpha 17$ to helix H7. The side wall formed by $\beta 4$ and $\beta 5$ is next to the active site, as indicated by lisinopril.


tACE: Natesh et al. Nature 421:551, 2003



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670

Fig. 2. Testicular ACE as redox- and mechanosensor.


tACE as redox sensor



Refs. Moskowitz et al, In press; Natesh et al, Nature 421:551, 2003


Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670

Fig. 3. *In silico* reduction of tACE: conversion of one cysteine to an alanine. "Reduction" of the middle cysteine of tACE removes a side wall of the active site. C352 and C370 in this drawing correspond to C383 and C401, respectively, in the numbering scheme of the tACE precursor (SwissProt P22966).



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670


Fig. 4a. Somatic ACE (sACE).



Title: Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s): Moskowitz
Appln. No. N/A
Docket # 44948/46670

Fig. 4b. Somatic ACE showing complementarity of the N- and C-terminal domains. The secretase, an integral membrane protein associated with sACE, may have a chaperone-like function to hold the two domains of sACE together.

Note: "Tongue-in-groove" fit



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670

Figure 5. The "disulfide zipper" at the heart of sACE: a potential electron transport chain.

sACE: Dimerization via a Disulfide Zipper





Figure 6. The disulfide isomerase exchange reaction possible at the heart of the disulfide zipper which might convert sACE into a kind of "reciprocating" enzyme.

sACE: a reciprocating enzyme with higher k_{cat} than tACE?




Activated by a single pair of electrons

S*, C-domain cysteine 341' or 360'


S*, N-DOMAIN CYSTEINE 341 OR 360

Figure 7. The two steps in the catalytic cycle of the "reciprocating" enzyme, sACE.




Title: Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s): Moskowitz
Appln. No. N/A
Docket # 44948/46670

Figure 9. NO inactivation of sACE: possible sites of S-nitrosylation.



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670


Figure 10. sACE on one immunocyte (T cell, monocyte/macrophage, or endothelial cell) binding to sACE on another immunocyte so as to promote specific cell-cell interactions. White triangle, "lid" formed by alpha helices $\alpha 1$ and $\alpha 2$; N and C, termini of each domain.



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670


Figure 11. Prompt response to angiotensin II receptor blockade in a 33 yr old white man with psoriasis. a,b,c: back and nail before treatment; d,e,f: 3 weeks after taking valsartan 80 mg daily. In a second case, a 62 yr old white man with chronic psoriasis, for which he took 75 mg methotrexate daily, was able to stop his methotrexate within 2 months of starting quinapril 200 mg/day.

a




b

Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670




c




d

Title: Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s): Moskowitz
Appln. No. N/A
Docket # 44948/46670




e

Title: Use of Angiotensin Receptor Blockers to Treat
Disease
Inventor(s): Moskowitz
Appl. No. N/A
Docket # 44948/46670



f

Title: Use of Angiotensin Receptor Blockers to Treat
Disease
Inventor(s): Moskowitz
Appl. No. N/A
Docket # 44948/46670




Title: Use of Angiotensin Receptor Blockers to Treat
Disease
Inventor(s): Moskowitz
Appln. No. N/A
Docket # 44948/46670

Figure 12. Genetic pathway for initiation of benign prostatic hyperplasia and prostate cancer. PDEF, prostate-derived ets factor.

Prostate Cancer in White Men

- ACE D/D is protective (& assoc'd w/ BPH)



Title:	Use of Angiotensin Receptor Blockers to Treat Disease
Inventor(s):	Moskowitz
Appln. No.	N/A
Docket #	44948/46670